



## Digital Receipt

This receipt acknowledges that **Turnitin** received your paper. Below you will find the receipt information regarding your submission.

The first page of your submissions is displayed below.

Submission author: Soni Andriawan  
Assignment title: Publication Articles Januari-Juni 2020  
Submission title: Study of *Holothuria scabra* effect on...  
File name: Soni\_Andriawan.pdf  
File size: 244.94K  
Page count: 10  
Word count: 5,358  
Character count: 30,555  
Submission date: 25-Feb-2020 03:43PM (UTC+0700)  
Submission ID: 1263783293



### Study of *Holothuria scabra* effect on the immune activity of *Pangasianodon hypophthalmus* against *Aeromonas hydrophila*

<sup>1</sup>Soni Andriawan, <sup>2</sup>Sofyan Hadi, <sup>3</sup>Sri Andayani, <sup>4</sup>Eilana Sanoesi, <sup>5</sup>Maftuch

<sup>1</sup> University of Muhammadiyah Malang, Indonesia; <sup>2</sup> PT. Mega Marine Pride, Pasuruan, Jawa Timur, Indonesia; <sup>3</sup> Brawijaya University, Malang, East Java, Indonesia.  
Corresponding author: S. Andriawan, soniandriawan@umm.ac.id

**Abstract.** *Holothuria scabra* or holothurians are one of the Echinodermata group which is specific and easily known. They have been known to be used as a drug to treat some diseases in humans. The aim of this study was to investigate the effect of *H. scabra* on the immune activity of *Pangasianodon hypophthalmus* by immersion and their protection against *Aeromonas hydrophila* invasion. In brief, *P. hypophthalmus* was immersed with *H. scabra* extract at 0, 50, 100 and 150 ppm by twice booster. Leucocyte level, number of phagocytes, phagocytosis, survival rate, and clinical symptoms were assessed to determine *P. hypophthalmus* immunity. The results revealed that the concentration of *H. scabra* extract at 100 ppm enhanced significantly the leucocyte level of *P. hypophthalmus* compared with the control. Concerning survival rate it was showed that all treatments worked better than the control. The highest percentage of survival rates was also observed at 100 ppm of *H. scabra* extract after challenged with *A. hydrophila*. The present discoveries suggest that *H. scabra* extract at the concentration of 100 ppm may raise the immune responses in *P. hypophthalmus* using immersion.

**Key Words:** holothurians, immersion, immune responses, iridescent shark, survival rate.

**Introduction.** Iridescent shark, *Pangasianodon hypophthalmus*, is one of the freshwater fish organisms having the fastest growth and potential economic value (Singh & Lakra 2012). In 2011, Indonesia was the second greatest *P. hypophthalmus* producer in the global manufacturing up to 16.1% of overall global production of *Pangasius*; however, the quantity is still below Vietnam which leads to world's *Pangasius* production amounting up to 80.9% (Ramadhan et al 2016). However, the intensive production of *P. hypophthalmus* with high density has posed a problem of infectious diseases (Afrianto & Liviawaty 1992). The most common disease-causing agents of *P. hypophthalmus* is bacteria (Griffiths et al 2010). In a recent study, *Aeromonas hydrophila* was observed infecting *P. hypophthalmus* (Griffiths et al 2010; Sarker & Faruk 2016). *A. hydrophila* is a gram-negative bacteria that cause motile *Aeromonas* septicemia in fish (Abdelhamed et al 2017). They produce exotoxins properties which cause hemorrhages on body, eyes, and fins, bloody ascites in the peritoneum, and contributing to swollen belly (Daskalov 2006; Griffiths et al 2010).

Today, antibiotics are still used to treat and prevent bacterial diseases (Iginosa et al 2012; Serano 2005). However, it is figured out that overuse of antibiotics can diminish the effectiveness of antibiotics (Sarum et al 2006), increase environmental hazards (Guo et al 2011) and generate residual antibiotics (Lakshmi et al 2013). One of the alternatives replaces antibiotics with natural products (Disch et al 2017). *Holothuria scabra* has been known to be used as a drug for treating several diseases in humans (Darsono 1993; Dhinakarana & Lipton 2014; Manan et al 2016). Contains compounds as steroid (Chalorak et al 2017; Goad et al 1985) Irterpenoid (Kerr & Chen 1995) saponin, alkaloid, glycosaminoglycan (Nagase et al 1995), lectins (Mojica & Merca 2005), phenol (Mameiona et al 2007) and flavonoid (Bordbar et al 2011; Nimah et al 2012). These compounds are used as an anticoagulant, antithrombotic, reducing cholesterol level and

# Study of *Holothuria scabra* effect on the immune activity of *Pangasianodon hypophthalmus* against *Aeromonas hydrophila*

*by* Soni Andriawan

---


**Submission date:** 25-Feb-2020 03:43PM (UTC+0700)

**Submission ID:** 1263783293

**File name:** Soni\_Andriawan.pdf (244.94K)

**Word count:** 5358

**Character count:** 30555



## Study of *Holothuria scabra* effect on the immune activity of *Pangasianodon hypophthalmus* against *Aeromonas hydrophila*

<sup>1</sup>Soni Andriawan, <sup>2</sup>Sofyan Hadi, <sup>3</sup>Sri Andayani, <sup>3</sup>Ellana Sanoesi, <sup>3</sup>Maftuch

<sup>1</sup> University of Muhammadiyah Malang, Indonesia; <sup>2</sup> PT. Mega Marine Pride, Pasuruan, Jawa Timur, Indonesia; <sup>3</sup> Brawijaya University, Malang, East Java, Indonesia.  
Corresponding author: S. Andriawan, soniandriawan@umm.ac.id

**Abstract.** *Holothuria scabra* or holothurians are one of the Echinodermata group which is specific and easily known. They have been known to be used as a drug to treat some diseases in humans. The aim of this study was to investigate the effect of *H. scabra* on the immune activity of *Pangasianodon hypophthalmus* by immersion and their protection against *Aeromonas hydrophila* invasion. In brief, *P. hypophthalmus* was immersed with *H. scabra* extract at 0, 50, 100 and 150 ppm by twice booster. Leucocyte level, number of phagocytes, phagocytosis, survival rate, and clinical symptoms were assessed to determine *P. hypophthalmus* immunity. The results revealed that the concentration of *H. scabra* extract at 100 ppm enhanced significantly the leucocyte level of *P. hypophthalmus* compared with the control. Concerning survival rate it was showed that all treatments worked better than the control. The highest percentage of survival rates was also observed at 100 ppm of *H. scabra* extract after challenged with *A. hydrophila*. The present discoveries suggest that *H. scabra* extract at the concentration of 100 ppm may raise the immune responses in *P. hypophthalmus* using immersion.

**Key Words:** holothurians, immersion, immune responses, iridescent shark, survival rate.

**Introduction.** Iridescent shark, *Pangasianodon hypophthalmus*, is one of the freshwater fish organisms having the fastest growth and potential economic value (Singh & Lakra 2012). In 2011, Indonesia was the second greatest *P. hypophthalmus* producer in the global manufacturing up to 16.1% of overall global production of Pangasius; however, the quantity is still below Vietnam which leads to world's Pangasius production amounting up to 80.9% (Ramadhan et al 2016). However, the intensive production of *P. hypophthalmus* with high density has posed a problem of infectious diseases (Afrianto & Liviawaty 1992). The most common disease-causing agents of *P. hypophthalmus* is bacteria (Griffiths et al 2010). In a recent study, *Aeromonas hydrophila* was observed infecting *P. hypophthalmus* (Griffiths et al 2010; Sarker & Faruk 2016). *A. hydrophila* is a gram-negative bacteria that cause motile Aeromonas septicemia in fish (Abdelhamed et al 2017). They produce exotoxins properties which cause hemorrhages on body, eyes, and fins, bloody ascites in the peritoneum, and contributing to swollen belly (Daskalov 2006; Griffiths et al 2010).

Today, antibiotics are still used to treat and prevent bacterial diseases (Igbinosa et al 2012; Serrano 2005). However, it is figured out that overuse of antibiotics can diminish the effectiveness of antibiotics (Sorum et al 2006), increase environmental hazards (Guo et al 2011) and generate residual antibiotics (Lakshmi et al 2013). One of the alternatives replaces antibiotics with natural products (Disch et al 2017). *Holothuria scabra* has been known to be used as a drug for treating several diseases in humans (Darsono 1993; Dhinakarana & Liptonb 2014; Manan et al 2016). Contains compounds as sterol (Chalorak et al 2017; Goad et al 1985) triterpenoid (Kerr & Chen 1995) sapogenin, alkaloid, glycosaminoglycan (Nagase et al 1995), lectins (Mojica & Merca 2005), phenol (Mamelona et al 2007) and flavonoid (Bordbar et al 2011; Nimah et al 2012). These compounds are used as an anticoagulant, antithrombotic, reducing cholesterol level and

blood fat, anticancer, antitumor, antibacterial, immun<sup>1</sup>stimulant, antifungal, antiviral, antimalarial and antiemetic (Bordbar et al 2011; Nimah et al 2012).

Therefore, the aim of this study was to examine leucocytes level, number of phagocytes, phagocytosis, survival rate and clinical symptoms of *P. hypophthalmus* and its tolerance to *A. hydrophila* post-immersion of *H. scabra* extract.

## Material and Method

**Fish and plant extraction.** *P. hypophthalmus* (6±1.0 g) were obtained from a commercial farm from Tulungagung, East java and reared in Parasite Laboratory, Faculty of Fishery and Marine Science<sup>1</sup> Brawijaya University, Malang, East java, Indonesia. Fish (10 individuals/aquarium) were acclimatized at 25°C for one week before conducting an experiment and fed with commercial feed at 3% of fish body weight per day. Water quality was controlled to keep the environment in good condition.

*H. scabra* extraction followed the study of Aras (2013) and Dong et al (2008) with slight modifications. Briefly, *H. scabra* was minced to small pieces, then 400 g of *H. scabra* was added to 400 mL of methanol and macerated in a bottle for 24 hours. The mixture was then filtrated to get methanol extract, while the waste was added again with methanol to isolate the extract completely. The clear supernatant of *H. scabra* extract was evaporated using vacuum evaporator to evaporate the solvent. Eight gram of thick extract was fractionated with water and n-butanol (1:1). The n-butanol fraction was evaporated using vacuum evaporator to obtain *H. scabra* extract. The extract was kept at 4°C until use.

**Effect of *H. scabra* extract on the immune level.** First, the *H. scabra* extract was dissolved directly into each aquarium with different concentrations of 0, 50, 100 and 150 ppm (T0, T1, T2, and T3, respectively) by twice booster at 0 and 7 day. The immersion of *H. scabra* extract was conducted for 1 hour and then *P. hypophthalmus* was transferred to each aquarium. The blood collection was conducted at 2, 4, and 6-day post-immersion of *H. scabra* extract and 3<sup>rd</sup> day post-infection to observe leucocyte level. Blood collection was taken from the caudal vena using sodium citrate 3% as anti-coagulant (Nuryati et al 2010).

Based on a method of Kamaludin (2011), leucocyte level was calculated under a microscope with 400x magnification. The number of leucocytes was formulated by (Suhermanto et al 2011):

$$SDP = (A/N) \times (1/V) \times Df$$

Where:

SDP = Number of leucocyte

A = Counted leucocyte

N = Number of hemocytometer square observed

V = Volume of hemocytometer square observed

Df = Dilution factor

The total number of macrophage cells was calculated based on the method of Irianto (2005). Fish kidney gently was powdered using tissue grinder and diluted in RPMI medium contained Fetal Bovine Serum (FBS) 2% in order to facilitate calculating the macrophage cells. Total macrophage cells was determined using a microscope with 40X magnification (Irianto 2005), while phagocytosis activity was determined following the Irianto & Austin (2002) method. Briefly, macrophage suspension was centrifuged with 1,000 rpm for 5 minutes and the pellet was spread on object glass. It was incubated at 26°C for 60 minutes and washed using RPMI medium contained FBS 2%. *A. hydrophila* was used for phagocytosis activity identification under a microscope with x100 magnification. The percentage of phagocytosis activity was calculated from 100 macrophage cells identified (Irianto & Austin 2002).

**Challenge and evaluation of survival rate.** The fish-pathogenic strain of *A. hydrophila* was kindly provided<sup>1</sup> by Microbiology Laboratory, Brawijaya University and used in this study with  $1 \times 10^8$  (CFU) mL<sup>-1</sup> for a challenge (LD 50). The challenge test was conducted at

day 8 of post-second booster (day 7). Ten fish were immersed with *A. hydrophila* ( $1 \times 10^8$  (CFU)  $m^{-1}$ ) into an aquarium. Survival rate of fish was monitored during 24, 48, 72, 96, and 144 hours post-challenge.

**Statistical analysis.** ANOVA (One-way analysis of variance) was applied to evaluate variations among groups. Multiple comparisons (BNT test) were carried out to determine significant differences among treatments using SPSS (version 17, USA). Data were displayed as the mean  $\pm$ SD,  $p < 0.05$  was considered significant.

## Results and Discussion

**Phytonutrients test.** Sea cucumber can produce highly active substances, saponins, the main secondary metabolites, which are the essential of their chemical defense (Zhao et al 2018). Based on Table 1, *H. scabra* extract was identified possessing compounds including saponin and triterpenoids. According to Zhao et al (2018), sea cucumber possesses some compounds which of the dominant compound are saponin (Table 1). Saponin is a glycoside triterpenoid obtained from secondary metabolite by membrane sterols which is linked by aglycone and sugar chains through the  $\beta$ -glycosidic bond (Bahrami et al 2014; Caulier et al 2016; Van Dyck et al 2011). This compound is known as a defense to predator, antimicrobial, and poison to invertebrate and algae (Van Dyck et al 2011).

Table 1  
Results of phytonutrients test

Test		Sign
Flavonoid	-	No color changed to red or orange
Saponin	✓	Foam formed
Steroid	-	Blue or purple color
Triterpenoid	✓	Red color
Fenol	-	No color changed to blue or green
Alkaloid	-	Brown sediment formed

- : negative expression; ✓ : positive expression.

**Immunological effect post-immersion.** The innate immune system is the first mechanisms defense of fish which is physiologically similar to greater vertebrates (Magnadóttir 2006; Uribe et al 2011). Their immune system is formed of physical barriers, cellular and humoral components which are used against foreign substances, such as microorganisms, toxins or malignant cells, responding to factors such as endogenous or exogenous components that stimulate this system (Biller-Takahashi & Urbinati 2014).

The inflammation is also recognized as an immune response, facilitated by complex interactions of cellular and humoral compounds (Biller-Takahashi & Urbinati 2014). The granulocytes are the initial cell that takes place at the inflammation site, being competent for eliminating pathogens. On the other hand, macrophage phagocytosis play an important role to remove the remaining pathogenic cells and cellular debris (Magnadóttir 2006).

**Leucocytes level.** Fish leucocytes could be used to determine fish health status or clinical evaluation of stress and diseases of fish (Palić et al 2011). According to Davis et al (2008), leukocyte studies are especially helpful to conservative physiology of fish because they are changed by stress and can be promptly connected to stress hormone levels. Leucocytes itself could be divided into three types such as monocytes, macrophages, and neutrophils (Mathias et al 2009; Neumann et al 2001). They play a role in intracellular and extracellular mechanisms to generate antimicrobial substances in defense against pathogens (Rieger & Barreda 2011).

In this study, the leucocytes level was enhanced post-immersion with *H. scabra* extract. The leucocytes level enhanced early 2<sup>th</sup>-day post-immersion by T1 and T2, while T3 showed no significant difference compared with the control group ( $p < 0.05$ ). Further, in another time point, leucocytes level was decreasing but it was still a significant difference comparing with control. In the 6<sup>th</sup> day, all treatments showed no significant differences, except T2 which still increased. Based on Suhermanto et al (2011) study, *H. scabra* extract also was applied for increasing leucocytes level of *Cyprinus carpio* pre-infection of *A. hydrophila*. Increased of leucocytes level was believed to be due to triterpenoids within *H. scabra* extract which have immune-regulatory function (Rascón-Valenzuela et al 2016). In another study, injection of triterpenoids enhanced white blood cells of mice ( $135.75 \pm 6.4\%$ ) at 3<sup>th</sup> and 6<sup>th</sup>-day post-injection (Raphael & Kuttan 2003). According to Aminin (2016), the small concentrations ( $0.1\text{--}6.0\text{ mg mL}^{-1}$ ) of holothurians triterpene glycosides play to trigger a leukocyte movement that promote an increase in phagocytosis of *Staphylococcus aureus* by human polymorphonuclear leucocyte. Based on it, we assumed that increasing of leucocytes was caused by triterpenoid compound within *H. scabra* extract (Figure 1).




Figure 1. Total leucocyte pre- and post-infection with *Aeromonas hydrophila*.

The second booster was conducted pre-challenge with *A. hydrophila* on 7<sup>th</sup> day. It was applied not only for increasing leucocyte again but also for showing the efficacy of *H. scabra* extract. In post-infection, the highest expression leucocytes level was showed by T0 and T3, while T1 and T2 showed no increase and dropped, respectively. In several cases, the leucocytes count increase post-infection of pathogen as a defense mechanism. According to Rajapakshe et al (2012) and Shahi et al (2014), an increase in the leucocyte level is the most reasonable results in an improvement of the non-specific defense, because macrophages and other phagocytic cells are main cells of the immune system. Interestingly, in our results, T1 and T2 did not enhance leukocytes level post-infection. Generally, a decreasing amount of leukocytes post-infection showed that leukocytes are believed active and come into blood vessel going through infected tissues (Nuryati et al 2010). Based on these findings, it is suggested that triterpenoids from *H. scabra* extract is recommendable to increase the immune response of *P. hypophthalmus*.

**Total of macrophage cells.** Macrophages can be simply collected from various sources such as blood (monocytes), lymphoid organs (specifically the kidney), and the peritoneal cavity (Secombes 1996). In this study, macrophages cells were isolated from kidney pre- and post-infection of *A. hydrophila*. The fish head kidney is believed equal of fish bone marrow-derived macrophages and has the capability of phagocytosis (Grayfer et al 2018), generate radicals (Grayfer & Belosevic 2009), and part of innate activation (Joerink et al 2006). Our results revealed that a total of macrophage cells increased significantly post-immersion with *H. scabra* extract and post-infection by all *H. scabra* concentrations. In pre-infection, T2 showed the most significant difference ( $p < 0.05$ )



compared with control and even with all treatments, while after infection all treatments showed significant differences ( $p < 0.05$ ) compared control group (Figure 2).




Figure 2. Total macrophage cells pre- and post-infection with *Aeromonas hydrophila*.

Increasing of macrophages post *H. scabra* extract immersion and *A. hydrophila* in *P. hypophthalmus* kidney could be possible due to leucocytes immigration. According to Grayfer et al (2018), Tissues macrophages were considered to result from circulating monocyte precursors in response to tissue entry and accompanying stimuli. An another study reported that phagocytic activity in tissues is facilitated by cytokines as macrophage stimulating factor released by peritoneal lymphocytes (Gordon & Plüddemann 2017; Graham & Secombes 1988). A similar result, reports that the use of 50 g *Euphorbia hirta* extract could enhance total macrophage in *C. carpio* kidney post- *E. hirta* feeding and *A. hydrophila* infection (Pratheepa & Sukumaran 2014). Based on our results, *H. scabra* extract is recommendable as immunostimulant to enhance total macrophage cells number in *P. hypophthalmus* kidney.

**Phagocytosis activity.** The fish kidney was considered as granulopoiesis site to produce granulocytes such as macrophages and neutrophils (Ainsworth 1992; Esteban et al 2015; Geven & Klaren 2017; Uribe et al 2011). They have been known as phagocytic cells that show a critical role in the fish nonspecific immunity, which have a phagocytosis activity (Hardi et al 2017; Romano et al 1998). Phagocytosis is typically a defensive reaction against invasion of foreign substances and, in the immune system, phagocytosis is a major mechanism used to remove pathogens and/or cell debris related function in engulfing and killing the pathogen (Esteban et al 2015).

In our results, phagocytic activity results showed significant differences among treatments ( $p < 0.05$ ) in *P. hypophthalmus* kidney by all concentrations post-immersion of *H. scabra* extract and even post-infection of *A. hydrophila* (Figure 3). Based on it, all concentrations of *H. scabra* extract worked in order to increase phagocytosis activity. According to (Hohn et al 2009), increasing of phagocytosis activity is considered by diverse factors such as pollutants (Weeks & Warinner 1986), diet (Blazer 1991), temperature (Hardie et al 1994), pathogens (Ainsworth & Dexiang 1990) and genetic variation (Sarder et al 2001). Phagocytic activity ratio (%) noted the highest value within T2 post-challenge with 41.89% activity followed by T1 and T3, respectively.

Aminin et al (2008) discovered that a small dosage of sea cucumber triterpenoid could enhance the antibody production to stimulate monocyte phagocytosis of *S. aureus* and the number of antibody such as IgM and IgG antibody levels in mice. An another study showed phagocytic activity increase in groups fed with 100 ng prolactin and 50 ng  $\beta$ -glucan per kg *Salmo salar* during the 20 days period (Paredes et al 2013). Based on it, we assumed that increase of phagocytic activity is enhanced by leucocytes migration from blood which goes through tissues.




Figure 3. Phagocytosis activity cells pre- and post-infection of *Aeromonas hydrophila*.

**Survival rate of *P. hypophthalmus*.** The survival rate was monitored for 144 hours post challenge with *A. hydrophila*. Challenge was conducted day 2 post-immersion of *H. scabra* extract (second booster) by bathing of *A. hydrophila*. All treatments showed better results than control ( $p < 0.05$ ). The highest expression of survival rate was recorded by T2 with 76.7%, while the lowest survival rate was shown by T0 with 16.7% (Table 2).

Table 2  
The survival rate of *Pangasianodon hypophthalmus* against *Aeromonas hydrophila*

Bacterial dose (cells/mL)	<i>H. scabra</i> extract (ppm)	No. of fish	Survival rate (%) at different times (hour)				
			24	48	72	96	144
No infection	0 (Control)	30	100	100	100	100	100
$10^8$	0	30	100	83.3	50	33.3	16.7
$10^8$	50	30	100	90	73.3	60	50
$10^8$	100	30	100	86.7	83.3	80	76.7
$10^8$	150	30	100	86.7	66.7	40	33.3

According to a study of Alipiah et al (2016), sea cucumber extract worked well to enhance survival rate of grouper post-challenge with several infectious bacteria. In a review, triterpenoid in *H. scabra* has a function as improvement of health status and can be used as a food supplement (Iñiguez-Martinez et al 2005). According to Nugroho et al (2016), triterpene was identified in *Terminalia catappa* which could enhance percentage of *Betta* sp. survival rate during cultivation. Based on the present study, *H. scabra* could improve health status of *P. hypophthalmus* post-immersion with *A. hydrophila*.

**Conclusions.** The results demonstrated that the immersion in *H. scabra* extract improved the survival rate of *P. hypophthalmus* after challenged with *A. hydrophila* and also enhanced the innate immunity of *P. hypophthalmus* by increasing the WBC, total macrophages, and phagocytosis activity that play crucial role in fish immunity.

**Acknowledgements.** The authors are grateful to Laboratory of Fish Parasite, Faculty of Fisheries and Marine Sciences, Brawijaya University, Indonesia. This research was supported by Advisor project and team.

## References

- Rascón-Valenzuela L. A., Torres Moreno H., Velazquez C., Garibay-Escobar A., Robles-Zepeda R. E., 2016 Triterpenoids: Synthesis, uses in cancer treatment and other biological activities. In: Advances in medicine and biology. Berhardt L. V. (ed), Chapter: 6.



- Abdelhamed H., Ibrahim I., Baumgartner W., Lawrence M. L., Karsi A., 2017 Characterization of histopathological and ultrastructural changes in channel catfish experimentally infected with virulent aeromonas hydrophila. Front Microbiol 8, 1519. doi:10.3389/fmicb.2017.01519.
- Afrianto I. E., Liviawaty I. E., 1992 Pengendalian hama & penyakit ikan: Kanisius.
- Ainsworth A. J., 1992 Fish granulocytes: Morphology, distribution, and function. Annual Review of Fish Diseases 2:123-148.
- Ainsworth A. J., Dexiang C., 1990 Differences in the phagocytosis of four bacteria by channel catfish neutrophils. Developmental & Comparative Immunology 14(2):201-209.
- Alipiah N. M., Ramli N. H. S., Low C.-F., Shamsudin M. N., Yusoff F. M., 2016 Protective effects of sea cucumber surface-associated bacteria against *Vibrio harveyi* in brown-marbled grouper fingerlings. Aquaculture Environment Interactions 8:147-155.
- Aminin D., 2016 Immunomodulatory properties of sea cucumber triterpene glycosides. In: Marine and freshwater toxins. Toxinology. Gopalakrishnakone P., Haddad Jr. V., Tubaro A., Kim E., Kem W. (eds), pp. 381-401, Springer, Dordrecht.
- Aminin D., Agafonova I., Kalinin V., Silchenko A., Avilov S., Stonik V., Collin P. D., Woodward C., 2008 Immunomodulatory properties of frondoside a, a major triterpene glycoside from the north atlantic commercially harvested sea cucumber cucumaria frondosa. Journal of Medicinal Food 11(3):443-453.
- Aras A., 2013 Uji toksisitas ekstrak teripang *Holothuria scabra* terhadap *Artemia salina*. Essay, Universitas Hasanuddin Makassar, Makassar.
- Bahrami Y., Zhang W., Chataway T., Franco C., 2014 Structural elucidation of novel saponins in the sea cucumber *Holothuria lessona*. Marine Drugs 12(8):4439-4473.
- Biller-Takahashi J. D., Urbinati E. C., 2014 Fish immunology. The modification and manipulation of the innate immune system: Brazilian studies. Anais da Academia Brasileira de Ciências 86(3):1484-1506.
- Blazer V. S., 1991 Piscine macrophage function and nutritional influences: A review. Journal of Aquatic Animal Health 3(2):77-86.
- Bordbar S., Anwar F., Saari N., 2011 High-value components and bioactives from sea cucumbers for functional foods - a review. Marine Drugs 9(10):1761-1805.
- Caulier G., Mezali K., Soualili D. L., Decroo C., Demeyer M., Eeckhaut I., Gerbaux P., Flammang P., 2016 Chemical characterization of saponins contained in the body wall and the Cuvierian tubules of the sea cucumber *Holothuria (Platyperona) sanctori* (Delle Chiaje, 1823). Biochemical Systematics and Ecology 68:119-127.
- Chalorak P., Jattujan P., Nobsathian S., Poomtong T., Sobhon P., Meemon K., 2017 *Holothuria scabra* extracts exhibit anti-parkinson potential in *C. elegans*: A model for anti-parkinson testing. Nutritional Neuroscience 21:1-12.
- Darsono P., 1993 Kandungan substansi bioaktif pada teripang oleh. ISSN.
- Daskalov H., 2006 The importance of *Aeromonas hydrophila* in food safety. Food Control 17(6):474-483.
- Davis A., Maney D., Maerz J., 2008 The use of leukocyte profiles to measure stress in vertebrates: A review for ecologists. Functional Ecology 22(5):760-772.
- Dhinakarana D. I., Liptonb A., 2014 Pharmacological potentials of sea cucumber *Holothuria atra* extracts from the Indian ocean. Asian Journal of Biomedical and Pharmaceutical Sciences 4(35), doi:https://doi.org/10.15272/ajbps.v4i35.537.
- Disch L., Drewe J., Fricker G., 2017 Dissolution testing of herbal medicines: Challenges and regulatory standards in europe, the united states, canada, and asia. Dissolution Technologies 24(2):6-12.
- Dong P., Xue C.-H., Du Q.-Z., 2008 Separation of two main triterpene glycosides from sea cucumber *Pearsonothuria graeffei* by high-speed countercurrent chromatography. Acta Chromatographica 20(2):269-276.
- Esteban M., Cuesta A., Chaves-Pozo E., Meseguer J., 2015 Phagocytosis in teleosts. Implications of the new cells involved. Biology 4(4):907-922.
- Geven E. J., Klaren P. H., 2017 The teleost head kidney: Integrating thyroid and immune signalling. Developmental & Comparative Immunology 66:73-83.

- Goad L., Garneau F.-X., Simard J.-L., ApSimon J., Girard M., 1985 Isolation of  $\Delta^{9(11)}$ -sterols from the sea cucumber *Psolus fabricii*. Implications for holothurin biosynthesis. *Tetrahedron Letters* 26(29):3513-3516.
- Gordon S., Plüddemann A., 2017 Tissue macrophages: Heterogeneity and functions. *BMC Biology* 15(1):53-53.
- Graham S., Secombes C. J., 1988 The production of a macrophage-activating factor from rainbow trout *Salmo gairdneri* leucocytes. *Immunology* 65(2):293-297.
- Grayfer L., Belosevic M., 2009 Molecular characterization, expression and functional analysis of goldfish (*Carassius auratus* L.) interferon gamma. *Developmental & Comparative Immunology* 33(2):235-246.
- Grayfer L., Kerimoglu B., Yaparla A., Hodgkinson J. W., Xie J., Belosevic M., 2018 Mechanisms of fish macrophage antimicrobial immunity. *Frontiers in Immunology* 9:1105. doi: 10.3389/fimmu.2018.01105.
- Griffiths D., Van Khanh P., Trong T., 2010 Cultured aquatic species information programme: *Pangasius hypophthalmus*. Fisheries and Aquaculture Department, Rome. Available at: [http://www.fao.org/fishery/culturedspecies/Pangasius\\_hypophthalmus/en](http://www.fao.org/fishery/culturedspecies/Pangasius_hypophthalmus/en). (Accessed 28 April 2015).
- Guo J. J., Her B. Y., Chou R. L., Chen T. I., 2011 Screening of modern herbal medicines in white shrimp (*Litopenaeus vannamei*) against *Vibrio harveyi* infection. *The Israeli Journal of Aquaculture*, Vol. 63, 7 p.
- Hardi E. H., Kusuma I. W., Suwinarti W., Saptiani G., Sumoharjo S., Lusiastuti A. M., 2017 Utilization of several herbal plant extracts on Nile tilapia in preventing *Aeromonas hydrophila* and *Pseudomonas* sp. bacterial infection. *Nusantara Bioscience* 9(2):220-228.
- Hardie L., Fletcher T., Secombes C., 1994 Effect of temperature on macrophage activation and the production of macrophage activating factor by rainbow trout (*Oncorhynchus mykiss*) leucocytes. *Developmental & Comparative Immunology* 18(1):57-66.
- Hohn C., Lee S.-R., Pinchuk L. M., Petrie-Hanson L., 2009 Zebrafish kidney phagocytes utilize macropinocytosis and  $Ca^{2+}$ -dependent endocytic mechanisms. *PLoS ONE*, 4(2), e4314-e4314. doi:https://doi.org/10.1371/journal.pone.0004314.
- Igbiosa I. H., Igumbor E. U., Aghdasi F., Tom M., Okoh A. I., 2012 Emerging aeromonas species infections and their significance in public health. *The Scientific World Journal* 2012:625023. doi: https://doi.org/10.1100/2012/625023.
- Iñiguez-Martínez A. M. d. M., Guerra-Rivas G., Ríos T., Quijano L., 2005 Triterpenoid oligoglycosides from the sea cucumber *Stichopus parvimensis*. *Journal of Natural Products* 68(11):1669-1673.
- Irianto A., 2005 Patologi ikan teleostei. Penerbit Gajah Mada University Press, Yogyakarta, Indonesia.
- Irianto A., Austin B., 2002 Use of probiotics to control furunculosis in rainbow trout, *Oncorhynchus mykiss* (Walbaum). *Journal of Fish Diseases* 25(6):333-342.
- Joerink M., Ribeiro C. M., Stet R. J., Hermesen T., Savelkoul H. F., Wiegertjes G. F., 2006 Head kidney-derived macrophages of common carp (*Cyprinus carpio* L.) show plasticity and functional polarization upon differential stimulation. *The Journal of Immunology* 177(1):61-69.
- Kamaludin I., 2011 Efektivitas ekstrak lidah buaya aloe vera untuk pengobatan infeksi *Aeromonas hydrophila* pada ikan lele dumbo *Clarias* sp. Melalui pakan.
- Kerr R. G., Chen Z., 1995 In vivo and in vitro biosynthesis of saponins in sea cucumbers. *Journal of Natural Products* 58(2):172-176.
- Lakshmi B., Viswanath B., Sai Gopal D., 2013 Probiotics as antiviral agents in shrimp aquaculture. *Journal of pathogens*. Article ID 424123, 13 pages <http://dx.doi.org/10.1155/2013/424123>.
- Magnadóttir B., 2006 Innate immunity of fish (overview). *Fish & Shellfish Immunology* 20(2):137-151.
- Mamelona J., Pelletier E., Girard-Lalancette K., Legault J., Karboune S., Kermasha S., 2007 Quantification of phenolic contents and antioxidant capacity of Atlantic sea cucumber, *Cucumaria frondosa*. *Food Chemistry* 104(3):1040-1047.

- Manan W. Z. W. A., Mahalingam S. R., Arshad K., Bukhari S. I., Ming L. C., 2016 Safety and efficacy of sea cucumber containing products. *Archives of Pharmacy Practice* 7:48-52.
- Mathias J. R., Dodd M. E., Walters K. B., Yoo S. K., Ranheim E. A., Huttenlocher A., 2009 Characterization of zebrafish larval inflammatory macrophages. *Developmental & Comparative Immunology* 33(11):1212-1217.
- Mojica E.-R. E., Merca F. E., 2005 Biological properties of lectin from sea cucumber (*Holothuria scabra* Jäger). *Journal of Biological Sciences* 5(4):472-477.
- Nagase H., Enjyoji K.-I., Minamiguchi K., Kitazato K. T., Kitazato K., Saito H., Kato H., 1995 Depolymerized holothurian glycosaminoglycan with novel anticoagulant actions: Antithrombin iii-and heparin cofactor ii-independent inhibition of factor x activation by factor ixa-factor viiia complex and heparin cofactor ii-dependent inhibition of thrombin. *Blood* 85(6):1527-1534.
- Neumann N. F., Stafford J. L., Barreda D., Ainsworth A. J., Belosevic M., 2001 Antimicrobial mechanisms of fish phagocytes and their role in host defense. *Developmental & Comparative Immunology* 25(8-9):807-825.
- Nimah S., Ma'ruf W. F., Trianto A., 2012 Uji bioaktivitas ekstrak teripang pasir (*Holothuria scabra*) terhadap bakteri *Pseudomonas aeruginosa* dan *Bacillus cereus*. *Jurnal Pengolahan dan Bioteknologi Hasil Perikanan* 1(1):9-17.
- Nugroho R., Manurung H., Saraswati D., Ladyescha D., M. Nur, F., 2016 The effects of *Terminalia catappa* L. Leaves extract on the water quality properties, survival and blood profile of ornamental fish (*Betta* sp.) cultured. *Biosaintifika* 8:241-248.
- Nuryati S., Maswan N., Alimuddin S., Sumantadinata K., Pasaribu F., Soejoeno R., Santika A., 2010 Gambaran darah ikan mas setelah divaksinasi dengan vaksin DNA dan diuji tantang dengan koi herpesvirus. *Jurnal Akuakultur Indonesia* 9(1):9-15.
- Palić D., Beck L. S., Palić J., Andreasen C. B., 2011 Use of rapid cytochemical staining to characterize fish blood granulocytes in species of special concern and determine potential for function testing. *Fish & Shellfish Immunology* 30(2):646-652.
- Paredes M., Gonzalez K., Figueroa J., Montiel-Eulefi E., 2013 Immunomodulatory effect of prolactin on atlantic salmon (*Salmo salar*) macrophage function. *Fish Physiology and Biochemistry* 39(5):1215-1221.
- Pratheepa V., Sukumaran N., 2014 Effect of euphorbia hirta plant leaf extract on immunostimulant response of aeromonas hydrophila infected *Cyprinus carpio*. *PeerJ*, 2, e671. doi:<https://doi.org/10.7717/peerj.671>.
- Rajapakshe A., Prasad K. P., Mukharjee S., 2012 Gross clinical signs and haematological changes associated with artificial infection of edwardsiella tarda in koi carp. *Journal of the National Aquatic Resources Research and Development Agency of Sri Lanka*, 41, pp. 1-10.
- Ramadhan A., Suwandi R., Trilaksani W., 2016 Competitiveveness strategies of indonesia pangasius fillet. *Indonesian Journal of Business and Entrepreneurship (IJBE)*, 2(2), 82. doi:<https://doi.org/10.17358/ijbe.2.2.82>.
- Raphael T. J., Kuttan G., 2003 Effect of naturally occurring triterpenoids glycyrrhizic acid, ursolic acid, oleanolic acid and nomilin on the immune system. *Phytomedicine* 10(6):483-489.
- Rieger A. M., Barreda D. R., 2011 Antimicrobial mechanisms of fish leukocytes. *Developmental & Comparative Immunology* 35(12):1238-1245.
- Romano N., Picchiatti S., Taverne-Thiele J. J., Taverne N., Abelli L., Mastrolia L., Verburg-van Kemenade B. M. L., Rombout J. H. W. M., 1998 Distribution of macrophages during fish development: An immunohistochemical study in carp (*Cyprinus carpio*, L.). *Anatomy and Embryology* 198(1):31-41.
- Sarder M. R. I., Thompson K. D., Penman D. J., McAndrew B. J., 2001 Immune responses of nile tilapia (*Oreochromis niloticus* L.) clones: I. Non-specific responses. *Developmental & Comparative Immunology* 25(1):37-46.
- Sarker J., Faruk M., 2016 Experimental infection of aeromonas hydrophila in pangasius. *Progressive Agriculture* 27(3), DOI: <https://doi.org/10.3329/pa.v27i3.30836>.
- Secombes C. J., 1996 2 - the nonspecific immune system: Cellular defenses. In: *Fish physiology*. Iwama G., Nakanishi T. (eds), Vol. 15, pp. 63-103, Academic Press.

- Serrano P. H., 2005 Responsible use of antibiotics in aquaculture. Food & Agriculture Organization.
- Shahi N., Mallik S., Sarma D., 2014 Leukocyte response and phagocytic activity in common carp, *Cyprinus carpio* experimentally infected with virulent *Aeromonas allosaccharophila*. Journal of Ecophysiology and Occupational Health 14(1-2):66-70.
- Singh A., Lakra W., 2012 Culture of *Pangasianodon hypophthalmus* into India: Impacts and present scenario. Pakistan Journal of Biological Sciences 15(1), 19. doi:10.3923/pjbs.2012.19.26
- Sørum M., Johnsen P., Aasnes B., Rosvoll T., Kruse H., Sundsfjord A., Simonsen G., 2006 Prevalence, persistence, and molecular characterization of glycopeptide-resistant enterococci in norwegian poultry and poultry farmers 3 to 8 years after the ban on avoparcin. Applied and Environmental Microbiology 72(1):516-521.
- Suhermanto A., Andayani S., Maftuch M., 2011 Pemberian total fenol teripang pasir (*Holothuria scabra*) untuk meningkatkan leukosit dan diferensial leukosit ikan mas (*Cyprinus carpio*) yang diinfeksi bakteri *Aeromonas hydrophila*. Jurnal Kelautan: Indonesian Journal of Marine Science and Technology 4(2):150-157.
- Uribe C., Folch H., Enriquez R., Moran G., 2011 Innate and adaptive immunity in teleost fish: A review. Veterinarni Medicina 56(10):486-503.
- Van Dyck S., Caulier G., Todesco M., Gerbaux P., Fournier I., Wisztorski M., Flammang P., 2011 The triterpene glycosides of *Holothuria forskali*: Usefulness and efficiency as a chemical defense mechanism against predatory fish. Journal of Experimental Biology 214(8):1347-1356.
- Weeks B., Warinner J., 1986 Functional evaluation of macrophages in fish from a polluted estuary. Veterinary Immunology and Immunopathology 12(1-4):313-320.
- Zhao Y. C., Xue C. H., Zhang T. T., Wang Y. M., 2018 Saponins from sea cucumber and their biological activities. Journal of Agricultural and Food Chemistry 66(28):7222-7237.

Received: 20 September 2019. Accepted: 12 December 2019. Published online: 19 December 2019.

Authors:

Soni Andriawan, University of Muhammadiyah Malang, Faculty of Agriculture and Animal Science, Department of Aquaculture, Indonesia, East Java, 65144 Malang, Jalan Raya Tlogomas 246, e-mail: soniandriawan@umm.ac.id

Sofyan Hadi, PT. Mega Marine Pride, Indonesia, Wonokoyo, Purwodadi, Gunungsari, Kec. Beji, Pasuruan, Jawa Timur, Postal Code: 67154, e-mail: sofyanhadi@gmail.com

Sri Andayani, University of Brawijaya, Faculty of Fishery and Marine Science, Department of Aquaculture, Indonesia, East Java, 65145 Malang, Jl. Veteran Malang, Ketawanggede, Kec. Lowokwaru, e-mail: yanik@ub.ac.id

Ellana Sanoesi, University of Brawijaya, Faculty of Fishery and Marine Science, Department of Aquaculture, Indonesia, East Java, 65145 Malang, Jl. Veteran Malang, Ketawanggede, Kec. Lowokwaru, e-mail: ellana.2011@ub.ac.id

Maftuch, University of Brawijaya, Faculty of Fishery and Marine Science, Department of Aquaculture, Indonesia, East Java, 65145 Malang, Jl. Veteran Malang, Ketawanggede, Kec. Lowokwaru, e-mail: maftuch@ub.ac.id

This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited.

How to cite this article:

Andriawan S., Hadi S., Andayani S., Sanoesi E., Maftuch, 2019 Study of *Holothuria scabra* effect on the immune activity of *Pangasianodon hypophthalmus* against *Aeromonas hydrophila*. AACL Bioflux 12(6):2167-2176.

# Study of Holothuria scabra effect on the immune activity of Pangasianodon hypophthalmus against Aeromonas hydrophila

## ORIGINALITY REPORT

7%

SIMILARITY INDEX

3%

INTERNET SOURCES

0%

PUBLICATIONS

9%

STUDENT PAPERS

## PRIMARY SOURCES

1

Submitted to Universitas Brawijaya

Student Paper

7%

Exclude quotes On

Exclude bibliography On

Exclude matches < 2%